

BIBLIOGRAPHY

- AGABA, P. 2002. Biological control of water hyacinth in the Kagera river headwaters of Ruwanda: A review through 2001. (Unpublished).
- CENTER, T. D. and VAN, T. K. 1989. Alternation of water hyacinth (*Eichhornia crassipes* {[Mart.]Solms} leaf dynamics and Phytochemistry by insect damage and plant density. Journal of Aquatic Botany, 35, 181-195.
- CENTER, T. D., ALLEN, F. D., JUBINSKY, G. P. and LESLIE, A.J. 1999. Water Hyacinth Weevils (*Neochetina eichhorniae* and *N. bruchi*) Inhibit Water Hyacinth (*Eichhornia crassipes*) Colony Development. Biological Control.15, 39-50.
- CENTER, T.D. and WRIGHT, N.1984. Biological control: Its place in the management of the Water hyacinth. Proceedings of the International Conference on Water Hyacinth (ed.G.Thyagarajan), 793-802. United Nations Environment Programme, Nairobi,Kenya.
- CENTER, T.D., HILL, M.P., CORDO, H. and JULIEN, M. H.2002. Water Hyacinth. Biological Control of Invasive Plants in the Eastern United States (ed. Driesche, V.). USDA Forest Service Publication 04, 413pp
- CHARUDATTAN, R. 2001. Biological Control of Water Hyacinth by Using Pathogens: opportunities, Challenges, and Recent Developments. ACIAR Proceedings 102, 21-28.
- CHIKWENHERE, G.P. 2001. Current strategies for the Mnagement of Water Hyacinth on the Manyame River System in Zimbabwe. ACIAR Proceedings 102, 33-38.
- DE GROOTE, H.,AJUONU, O., ATTIGNON, S.,DJESSOU,R. and NEUENSCHWANDER, P. 2002. Economical Economics. 45105-117.

DELOACH, C.J. and CORDO, H.A.1976. Ecological studies of *Neochetina eichhorniae* and *N.bruchi* on water hyacinth in Argentina. Journal of Aquatic Plant Management, 14, 53-59.

DELOACH, C.J. and CORDO, H.A.1976. Ecological studies of *Neochetina eichhorniae* and *N.bruchi* on water hyacinth in Argentina. Journal of Aquatic Plant Management, 14, 53-59.

DENOTH, M., FRID,L. and MYERS, J. H.2001.Multiple agents in biological control: improving the odds?. Biological Control 24,20-30.

FAYAD, Y.H., IBRAHIM, A.A., EL-ZOGHBY, A.A. and SHALABY, F.F. 2001. Ongoing Activities in the Biological Control of Water Hyacinth in Egypt. ACIAR Proceedings 102, 43-46.

FERNANDO, I.V.S. 1996.  University of Moratuwa Sri Lanka Electronic Theses & Dissertations www.lib.mrt.ac.lk Biological control of aquatic weeds in Sri Lanka. Proc. Sri Lanka Ass. Advmt. Sci. 52(2). 73-79.

GOPAL, B. 1987. Aquatic plant studies 1:water hyacinth. Elsevier Science Publishers, 5-161.

GRODOWITZ, M., CENTER, T. D. and FREEDMAN, J. E.1997. A Physiological Age – Grading System for *Neochetina eichhorniae* (Warner) (Coleoptera: Curculionidae), a Biological Control Agent of Water Hyacinth, *Eichhornia crassipes* (Mart.) Solms. Biological Control 9,89-105.

HEARD, T. A. and WINTERON, S. L. 2000. Interactions between nutrient status and weevil herbivory in the biological control of water hyacinth. Journal of Applied Ecology, 37, 117-127.

HILL, G., PHIRI, G., LWANDA, C., NJAYA, F., CHIMATIRO, S. and HILL, M.P. 1997. Water hyacinth Biological Control in the Shire River, Malawi. Proc. 1st IOBC Water hyacinth Working Group. 39-50.

HILL, M. P. and OLCKERS, T. 2001. Biological control initiatives against Water hyacinth in South Africa: Constructing factors, success and new courses. ACIAR Proceedings 102, 33-38.

JIANQING, D., REN, V., WEIDONG, F. and GUOLIANG, Z. 2001. Water Hyacinth in China: Its distribution, Problems and Control status. ACIAR Proceedings 102, 29-38.

JULIEN, M. H. 2001. Biological Control of Water Hyacinth with Arthropods. ACIAR Proceedings 102, 08-20.

JULIEN, M. H., GRIFTHS, M. W. and WRIGHT, A. D. 1999. Biological Control of Water hyacinth. The weevils, *Neochetina bruchi* and *N.eichhorniae*: biologies, host ranges, and rearing ,releasing and monitoring techniques for biological control of *Eichhornia crassipes*. ACIAR. Canberra.

KANNAN, C. and KATHiresan, R. M. 1997. Biological control at different growth stages of water hyacinth. Proc. 1st IOBC Water hyacinth Working Group.

KONDO, T. and SATO, H. 1981. Biomass production of water hyacinth and its ability to remove minerals from water: effect of the concentration of culture solution on the rates of plant growth and nutrient uptake. Japan Journal of Ecology, 31, 257-267.

MAKHANU, K. S. 1997. Impact of Water hyacinth on Lake Victoria. 23rd WEDC Conference, South Africa. 165-166

MOORHOUSE, T.M., AGABA, P. and MCNABB, T. J. 2001. Recent Efforts in Biological Control of Water Hyacinth in the Kagera River Headwaters of Ruwanda. ACIAR Proceedings 102, 39-42.

PANTILU, V.R.1984. Water Resource Development and Aquatic Weed Management. Proceedings of the International Conference on Water Hyacinth (ed.G.Thyagarajan), 737-749. United Nations Environment Programme, Nairobi,Kenya.

PENFOUND, T. and EARLE, T.T. 1948.The Biology of Water hyacinth. Ecological Monographs 18(4). 447-472.

ROOM, P. M. and FERNANDO, I.V.S.1992.Weed invasions countered by biological control: *Salvinia molesta* and *Eichhornia crassipes* in Sri Lanka. Journal of Aquatic Botany, 42, 99-107.

SALISBURY, F. B. and C. W. ROSS, 1990. Plant Physiology. Wadsworth Publishing Company, Belmont. CA.

WILSON, J.R., REES, M., HOLST, N., THOMAS, M. B. and HILL, G. 2001. Water hyacinth Population dynamics. ACIAR Proceedings 102, 33-38.

WRIGHT, A.D.1984. Effect of Biological Control Agents on Water Hyacinth in Australia. Proceedings of the International Conference on Water Hyacinth (ed.G.Thyagarajan), 823-833. United Nations Environment Programme, Nairobi, Kenya.