


LB 10005/20/2011

135

LIBRARY
UNIVERSITY OF MORATUWA, SRI LANKA
MORATUWA

IDENTIFICATION OF FACTORS FOR PREVALENCE OF OBESITY IN SRI LANKA



H D D G Henadheera

(8118)

Dissertation submitted in partial fulfillment of the requirements for the
degree Master of Operational Research

Department of Mathematics

University of Moratuwa
Sri Lanka

September 2010

96818

51 "10"
59.8 (043)




96818



DECLARATION


I declare that this is my own work and this dissertation does not incorporate without acknowledgement any material previously submitted for a Degree or Diploma in any University or other institute of higher learning and to the best of my knowledge and belief it does not contain any material previously published or written by another person except where the acknowledgement is made in the text.

Signature: 

Date: 16/12/2010

Copyright Statement


I hereby grant the University of Moratuwa the right to archive and to make available my dissertation in whole or part in the University Libraries in all forms of media, subject to the provisions of the current copyright act of Sri Lanka. I retain all proprietary rights, such as patent rights. I also retain the right to use in future works (such as articles or books) all or part of this dissertation.

Signature: 

Date: 16/12/2010

I have supervised and accepted this dissertation for the award of the degree.

Signature of the supervisor:



Date: 16/12/2010

ACKNOWLEDGEMENT

To My Beloved
Amma & Thaththa

ACKNOWLEDGEMENT

It is with deepest respect and gratitude I express my heartfelt thanks to my supervisor Dr M D T Attygalle, Head of the Department, Department of Statistics, University of Colombo who unremittingly leaned me an open ear, providing guidance and encouragement throughout to make this project a great success.

I am thankful to, Dr Shanthi Gunawardana, Nutrition Coordination Unit, Dr C D Gunarathne, Director, Nutrition Coordination Unit, Mr D Munidasa, Head, Computer Division, Nutrition Coordination Unit, and special thanks to Mrs. Mala Nandani Rajapakha, Director Administration (II), Department of Health Services for providing me with secondary data and background information for this project.

I like to take this opportunity to thank Dr. T.S.G. Peris, Coordinator, M.Sc. in Operational Research and Mr. T.M.J.A. Cooray former Coordinator, M.Sc. in Operational Research for providing me knowledge and guidance to carry out this project and all the members of the staff for providing help whenever needed.

I dearly express my appreciation and a big thank you to all my friends who willingly supported me to make this project a success. I am grateful to my dearest sister and brother who provided me a helping hand whenever needed and to my beloved parents for their valuable advice, encouragement and unconditional assistance given to me without which this report would not have been completed in this manner.

Abstract

Obesity is defined as an excessively high amount of body fat or adipose tissue in relation to lean body mass. This condition occurs as a result of higher energy intake than the energy expenditure. The Body Mass Index (BMI) is the tool utilized for the identification of obesity. A BMI greater than or equal to twenty five and less than thirty is considered as 'overweight' and a BMI greater than or equal to thirty is considered as 'obese'. Obesity is prevalent among people belonging to any layer of socio economic group or age group. The rate of increment of obesity prevalence has increased over the past few decades. The main causes of this have been identified as the environmental factor, the change in life style of people with the modernization and globalization trends and also genetics according to research experts. Obesity is revealed to be associated with many chronic diseases such as type 2 diabetes, cardiovascular disease, stroke and hypertension to name a few. In Sri Lanka obesity is being subject to much research and the Nutrition Coordination Department (NCD) has conducted a survey on prevalence of obesity island wide excluding North and East. NCD has analyzed the data at a descriptive level and the data set collected in the survey was used in this study for statistical analysis. The main objectives of this study were twofold. The first was to identify the possible risk factors for BMI based weight category of a respondent. The second was to statistically explore how the identified risk factors were related with BMI based weight category of a respondent.

A descriptive analysis and a univariate analysis were done followed by the main component, statistical modeling with an approach based on a proportional odds model. It was aimed to explore how the explanatory variables were associated with the BMI based weight category of a respondent which was the response variable under consideration. A recently developed accurate method of checking the goodness of fit of the proportional odds model was also used to ensure the adequacy of the model and the validity of the results of the analysis. The study focused for possible risk factors for prevalence of obesity in Sri Lanka with respect to respondent's socio-demographic information, respondent's history of diseases, family history of relevant diseases, smoking history, alcohol consumption and physical activity. As such there were thirty eight explanatory variables in these five areas and it was found that twenty two of them were significantly associated with the response variable which was BMI based weight category of the respondent. Fitting the proportional odds model revealed that BMI based weight category of a respondent is caused due to occupation, smoking, age, hypertension, walking, alcohol consumption, physical exercises and mother's diagnosis of Diabetes Mellitus. As such, physical activity in working and domestic life showed lesser probability of being overweight or obese. Alcohol consumers had a higher probability of being overweight or obese where as smokers were found to have an inverse relationship with BMI based weight category. Respondents who were diagnosed with Hypertension had a higher probability of being overweight or obese. Moreover people in the age group of 30-50 years were found to have the highest probability of being overweight or obese. Further it was found that a respondent is more vulnerable to be overweight or obese if the mother has been diagnosed with Diabetes Mellitus.

In conclusion, occupation, smoking, age, hypertension, walking, alcohol consumption, physical exercises and mother's diagnosis of Diabetes Mellitus was identified as the statistically significant risk factors for the prevalence of obesity in Sri Lanka. These statistically proven findings of this study are constructive of developing strategies and health policies in Sri Lanka for prevention of obesity, so as to ensure a healthier life style and to improve the quality of life of Sri Lankans.

TABLE OF CONTENTS

Declaration		i
Dedication		ii
Acknowledgement		iii
Abstract		iv
Table of Contents		v
List of Figures		vii
List of Tables		ix
List of Appendices		x
Chapter 1:	Introduction	1
	1.1 Background of the Study	1
	1.1.1 About the survey	4
	1.1.2 Sample Design	5
	1.2 Objectives of the Study	6
	1.3 Significance of the Study	6
	1.4 Statistical Methods and Preview of the Dissertation	6
Chapter 2:	Literature Review	7
Chapter 3:	Theory and Methodology	13
	3.1 Descriptive Analysis	13
	3.2 Univariate Analysis	13
	3.2.1 Parametric test	13
	3.2.2 Non-parametric test	15
	3.3 Logistic Regression Modeling	16
	3.4 Polytomous Response and Multiple Response Variables	16
	3.4.1 Proportional odds model	17
	3.5 Model Fitting	19
	3.5.1 Maximum Likelihood Method	19
	3.5.2 Parameter Estimates	19
	3.5.3 Solving Likelihood Equation	20

3.5.4	Comparing Alternative Logistic Models	20
3.5.5	Model Building	21
3.6	Odds Ratios	22
3.6.1	Odds Ratios for categorical explanatory variables when the variables are not involved in an interaction term	22
3.6.2	Odds Ratios for categorical explanatory variables when the variables are involved in an interaction Term	23
3.7	Goodness of fit of the model	25
3.7.1	Pearson Chi Square Statistic Deviance Statistic	25
3.7.2	Alternative Method in testing Goodness of fit	26
Chapter 4:	Preliminary Analysis	30
4.1	Descriptive Analysis	30
4.2	Univariate Analysis	50
Chapter 5:	Proportional Odds Model	52
5.1	Selected Variables	52
5.2	Proportional odds model	54
5.3.	Model Diagnostics	74
5.3.1	Score test	74
5.3.2	Goodness of fit tests	74
5.3.3	Alternative Method	75
5.3.4	Expected Counts and Residuals	78
5.3.5	Plot of Standardized Residuals of the Proportional Odds Model	81
5.4	Parameter estimates	82
5.5	Interpretation of odds ratios	83
Chapter 6:	Discussion	87
References		97
Appendix A:	Fitting second two factor interaction term	99
Appendix B:	SAS Code	105
Appendix C:	Questionnaire	109



LIST OF FIGURES

	Page
Figure 1.1: Graph of body mass index	4
Figure 4.1: Pie chart for body mass index	30
Figure 4.2: Pie chart for gender	31
Figure 4.3: Stacked column chart for BMI by gender	32
Figure 4.4: Stacked column chart for BMI by age group	32
Figure 4.5: Stacked column chart for BMI by marital status	33
Figure 4.6: Stacked column chart for BMI by number of children	33
Figure 4.7: Stacked column chart for BMI by educational level	34
Figure 4.8: Stacked column chart for BMI by occupation	35
Figure 4.9: Pie chart for religion	36
Figure 4.10: Stacked column chart for BMI by religion	37
Figure 4.11: Stacked column chart for BMI by nature of monthly income	38
Figure 4.12: Stacked column chart for BMI by history of respondent's illnesses	39
Figure 4.13: Stacked column chart for BMI by treatment for respondent's illnesses	40
Figure 4.14: Stacked column chart for BMI by medicine for other illnesses	40
Figure 4.15: Stacked column chart for BMI by if respondent ever smoked	41
Figure 4.16: Stacked column chart for BMI by if respondent's spouse smoked	42
Figure 4.17: Stacked column chart for BMI by if respondent ever consumed alcohol	42
Figure 4.18: Stacked column chart for BMI by if respondent walk to meet daily needs	43
Figure 4.19: Stacked column chart for BMI by respondent's number of sitting hours per day	44
Figure 4.20: Stacked column chart for BMI by respondent's number of sleeping hours per day	44

	Page
Figure 4.21: Stacked column chart for BMI by respondent's occupation involves manual work	45
Figure 4.22: Stacked column chart for BMI by if respondent do regular exercise	46
Figure 4.23: Stacked column chart for BMI by respondent's father's illnesses	47
Figure 4.24: Stacked column chart for BMI by respondent's mother's illnesses	48
Figure 4.25: Stacked column chart for BMI by respondent's sibling's illnesses	49
Figure 5.1: Plot of standardized residuals vs index group_response level	81
Figure 5.2: Plot of R^* vs index group_response level	81

LIST OF TABLES

	Page
Table 3.1: Two way contingency table	13
Table 3.2: Table for the two variables	15
Table 4.1: Comparison of sample and population figures by sex	31
Table 4.2: Comparison of sample and population figures by religion	36
Table 4.3: Results of the univariate analysis	50
Table 5.1: Variable description	52
Table 5.2: Results of fitting null model	54
Table 5.3: Results of fitting main effects to model 01	54
Table 5.4: Results of fitting main effects to model 02	56
Table 5.5: Results of fitting main effects to model 03	57
Table 5.6: Results of fitting main effects to model 04	58
Table 5.7: Results of fitting main effects to model 05	59
Table 5.8: Results of fitting main effects to model 06	60
Table 5.9: Results of fitting main effects to model 07	61
Table 5.10: Results of fitting main effects to model 08	62
Table 5.11: Results of fitting main effects to model 09	63
Table 5.12: Results of fitting second order interaction term to model 10	64
Table 5.13: Results of fitting second order interaction term to model 11	99
Table 5.14: Comparison of the results of the Score test	73
Table 5.15: Parameter estimates of the best model	73
Table 5.16: Results of the score test	74
Table 5.17: Results of goodness of fit test	75
Table 5.18: Results of goodness of fit test for alternative model	77
Table 5.19: Results of residual analysis for the proportional odds model	79
Table 5.20: Parameter estimates	82
Table 5.21: Odds ratios and confidence intervals	83

LIST OF APPENDICES

Appendix	Description	Page
Appendix - A	Fitting second two factor interaction term	99
Appendix – B	SAS Code	105
Appendix – C	Questionnaire	109