LB/DON/86/86

# Conservation of Masbedda (Gymnema sylvestre R. Br.) through propagation

py

## K. K. I. U. A. Kumara

## UNIVERSITY OF MORATUWA, SRI LANKA

This thesis was submitted to the Department of Civil Engineering of the University of Moratuwa in partial fulfilment of the requirements for the Degree of Master of Science in Environmental Management

504 (043)

Department of Civil Engineering
University of Moratuwa
Sri Lanka
August, 2004

85399

University of Moratuwa

#### **DECLARATION**

The work included in the thesis in part or whole has not been submitted for any other academic qualification at any institution.

Date: 02/08/2004

K. K. I. U. A. Kumara

I certify that the above statement is correct.

Date: 02 4 2004

**UOM Verified Signature** 

Dr. S. Subasinghe Senior Lecturer Department Crop Science

Faculty of Agriculture



Date: 4 /08/2004

**UOM Verified Signature** 

Dr. Maresh Jayaweera

Senior Lecturer

Department of Civil Engineering

University of Moratuwa

Moratuwa

#### **ABSTRACT**

\* 15'4 "

Several activities could be undertaken aimed at conservation and sustainable use of medicinal plants. Some of them are undertaken directly at locations where plants are naturally occurring, while others are less direct, such as commercial cultivation systems. Cultivation of medicinal plants is particularly important for species, which already have a good market demand. Gymnema sylvestre is one such rare medicinal plant. Therefore, the present study was aimed at developing cost effective and simple propagation techniques for Gymnema sylvestre.

Investigations on both sexual and vegetative means of propagation of *Gymnema* have been carried out, at the Faculty of Agriculture, University of Ruhuna, Mapalana, Kamburupitiya from October 2002 to November 2003. Viable, uniform seeds were screened and used for seed propagation studies. The effect of sowing media on germination of *Gymnema* seeds has been investigated using four kinds of sowing media, sand, coir dust, a mixture of top soil and sand (1:1) and a mixture of sand top soil and coir dust (1:1:1). Seeds were stored in an open system under room temperature (30 °C) and germination percentage was recorded at 15-day intervals. This was to find out the effect of storage period on germination of seeds. Complete Randomized Design was used for the experiments with five replicates, each contained ten seeds. Germination percentage was calculated and the experiments were repeated to confirm the results.

For vegetative propagation studies, three separate experiments were carried out using stem cuttings. Double nodal semi-hard wood cuttings were rooted in polybags filled with different rooting media including sand, a mixture of sand and top soil (1:1), a mixture of sand, top soil and compost (1:1:1) and top soil alone to investigate the effect of media on rooting. Hard wood, semi-hard wood and soft wood cuttings were planted in polybags filled with a mixture of sand, top soil and compost (1:1:1), to determine the effect of maturity of cuttings on rooting. Effect of watering on rooting of cuttings was investigated using semi-hard wood cuttings planted in polybags contained a potting mixture of sand, top soil and compost (1:1:1) with three watering frequencies. Complete Randomized Design was used for the studies with ten replicates. Number of roots, root length and root biomass accumulation were measured at the end of 6,10 and 14 weeks after planting.

Results showed that germination percentage of seeds was significantly ( $p \le 0.05$ ) high (92 %) in coir dust, whereas the lowest germination percentage (28 %) was observed from top soil media. There were no significant ( $p \le 0.05$ ) differences in germination of seeds throughout the first two months of storage under normal condition. Results of the vegetative propagation studies revealed that a mixture of sand, top soil and compost (1:1:1) was the most suitable rooting media for *Gymnema* cuttings, whereas the semi-hard wood cuttings rooted significantly ( $p \le 0.05$ ) higher than the hard wood and soft wood cuttings. With regard to the watering frequency, cuttings watered once in two days rooted and performed significantly ( $p \le 0.05$ ) better than the other treatments.

\* 04 % T

Results can be concluded that *Gymnema sylvestre* can be propagated by means of both sexual and asexual methods. Semi-hard wood cuttings, which appear to be more amenable to rooting and potting mixture of sand, top soil, and compost watered once in two days, showed the best results, while high germination percentage of seeds could be obtained within first two months of storage.



#### **ACKNOWLEDGEMENTS**

. v.d ..

I wish to express my deep gratitude and sincere appreciation to my supervisor Dr. S. Subasinghe, Senior Lecturer/Head, Department of Crop Science, Faculty of Agriculture, University of Ruhuna, for the encouragement and guidance; continued support and invaluable advice given to me throughout this study and allowing me to use facilities in the Department of Crop Science, in order to make this study a success.

I sincerely thank Dr. Mahesh Jayaweera, Course Co-ordinator, M.Sc./P.G. Diploma in Environmental Engineering and Management, for making arrangements to conduct this study at the Faculty of Agriculture, University of Ruhuna, and for his continued supervision and invaluable advice.

My thanks are due to Prof. (Mrs) N. Ratnayeke, Director, Environmental Division, Department of Civil Engineering, University of Moratuwa, for spending her valuable time with us, throughout the course and continued encouragement.

I would also like to thank Dr. M. de S. Liyanage, Senior Lecturer, Department of Crop Science, Faculty of Agriculture, University of Ruhuna, for his valuable advice and guidance.

University of Moratuwa, Sri Lanka.

I very much like to thank all the non-academic staff in the Environmental Division of the Department of Civil Engineering, University of Moratuwa for their assistance throughout the study period.

I am sincerely grateful to the staff of the Department of Crop Science, especially to Miss Dilusha B. Weerathunge for her assistance in data analysis.

Last, but not least, I take this opportunity to thank the Sri Lanka Conservation and Sustainable Use of Medicinal Plants Project for their financial assistance to support this study.

## LIST OF CONTENTS

		Page
ΑB	STRACT	I
AC	ACKNOWLEDGEMENTS	
LI	LIST OF CONTENTS	
LĽ	ST OF TABLES	VII
LĽ	ST OF FIGURES	VIII
LI	IST OF PLATES	
AF	BBREVIATIONS	X
1.	INTRODUCTION	1
	1.1 General introduction	1
	1.2. Objectives of the study	4
	University of Moratuwa, Sri Lanka. Electronic Theses & Dissertations www.lib.mrt.ac.lk	
2.	LITERATURE REVIEW	5
	2.1.Global use of medicinal plants	5
	2.2. Value of medicinal plants	6
	2.3. Loss of biological diversity and the availability of medicinal plants	8
	2.4. Future demand for medicinal plants	9
	2.5. Conservation of medicinal plants	10
	2.6. Cultivation of medicinal plants	11
	2.7. Use of medicinal plants in Sri Lanka	12
	2.8. General introduction to Gymnema sylvestre	14
	2.8.1. Botanical description	14
	2.8.2. Distribution	15
	2.8.3. Historical use	15
	2.8.4. Present uses	15
	2.9. Plant propagation	15
	2.9.1. Vegetative propagation	16
	2.9.1.1. Propagation of plants through cuttings	16

2.9.1.2. Stem cuttings	16
2.9.1.3. Soft wood cuttings	16
2.9.1.4. Semi- hard wood cuttings	17
2.9.1.5. Hard wood cuttings	17
2.9.1.6. Rooting media	17
2.9.1.7. Stock plants	18
2.9.2. Seed propagation	18
2.9.2.1. Sign of maturity	18
2.9.2.2. Factors influencing storage	18
2.9.2.3. Medium for germination	19
2.9.2.4. Importance of moisture	19
3. MATERIALS AND METHODS	20
3.1. Site description	20
3.2. Seed propagation	20
3.2.1. Seed collection Moralina, Sri Lanka.	20
3.2.2. Seed cleaning means the second	20
3.2.3. General information about seeds	22
3.2.4. Germination tests	22
3.2.5. Experiment 1	22
3.2.6. Experiment 2	23
3.2.7. Experiment 3	23
3.3. Vegetative propagation	23
3.3.1. Preparation of cuttings	23
3.3.2. Placing of the cuttings	24
3.3.3. Environmental modification	24
3.3.4. Experiment 4	24
3.3.5. Experiment 5	25
3.3.6. Experiment 6	25
3.4. Data analysis	26

\* 000 V .

4.	RESULTS AND DISCUSSION	27
	4.1. General information about seeds	27
	4.2. Seed propagation	29
	4.3. Vegetative propagation	35
5.	CONCLUSIONS	47
6.	FUTURE RECOMMENDATIONS	48
7.	REFERENCES	49



## LIST OF TABLES

	Page
Table 1 Number and percentage of medicinal plants used by countries	6



## LIST OF FIGURES

	Page
Figure 4.1 Effect of sowing media on germination of Gymnema seeds	30
Figure 4.2 Effect of moisture on germination of Gymnema seeds	31
Figure 4.3 Effect of storage period on germination of Gymnema seeds	32
Figure 4.4 Effect of cutting type on number of roots	35
Figure 4.5 Effect of cutting type on root length	36
Figure 4.6 Effect of cutting type on root biomass	37
Figure 4.7 Effect of media on number of roots	39
Figure 4.8 Effect of media on root length	40
Figure 4.9 Effect of media on root biomass rative. Sri Lanka.	41
Figure 4.10 Effect of watering frequency on number of roots	43
Figure 4.11 Effect of watering frequency on root length	44
Figure 4.12 Effect of watering frequency on root biomass	45



#### LIST OF PLATES

	Page
Plate 3.1. Mother plant stock	21
Plate 4.1. Gymnema plants at flowering stage	27
Plate 4.2. Gymnema pods	28
Plate 4.3. Gymnema seeds	29
Plate 4.4. Gymnema seedlings	34
Plate 4.5. Gymnema rooted cuttings	38
Plate 4.6. Gumnoma vegetatively propagated plants	42



#### **ABBREVIATIONS**

MAP - Medicinal and Aromatic Plants

WHO – World Health Organization

WL<sub>2</sub> - Low Country Wet Zone

CRD - Completely Randomized Design

DMRT - Duncan's Multiple Range Test

WAP - Weeks After Planting

