

REFERRANCEES

- [1] J. Perry, *Gait analysis, normal and pathological function*, 1 st ed., NJ:Slack, 1992, pp. 1-47, 111-129.
- [2] The American Heritage® Medical Dictionary Copyright © 2007, 2004 by Houghton Mifflin Company.
- [3] Andriacchi, T.P., Alexander, E.J., et. al ., 1998, "A Point Cluster Method for In Vivo Motion Analysis: Applied to a Study of Knee Kinematics", *J. Biomech Eng*, 120(12) p743-749,
- [4] Katherine F. Wells, Ph.D "Kinesiology" 1992:7:418-420.
- [5] Rockar,P.A. "The subtalar joint :Anatomy and Joint motion". *Journal of Orthopedic and sports physical therapy*.1995 21, p361-372.
- [6] Leardini A, Benedetti MG, Catani F, Simoncini L, Giannini S. "An anatomically based protocol for the description of foot segment kinematics during gait". *Clinical Biomechanics*. 1999;14:528-36.
- [7] Vaughan, et. a l . "Dynamics of human gait, Champaign Illinois": Human Kinetics Publishers
- [8] Murray,M.P.,A.B.Drought,and R.C.Kory, *walking pattern of normal men. Bone and Joint Surgery*,1964. 46(A);p 355-360.
- [9] Peter K. Larsen, 2Eva N. G. Pedersen, 3Tine Alkjær, 4Bente R. Jensen and 3Erik B. Simonsen; "Test-retest reliability of a video-based gait analysis system".
- [10] T.P. Andriacchi, and D.E. Hurwitz, "Gait biomechanics and the evolution of total joint replacement," *Gait and Posture*, vol. 5, pp. 256-264, 1997.
- [11] Kaltenborn.F.M "Manual Mobilization of the Extremity Joints" (4th ed.).p.27Minneapolis:OTTP.
- [12] Martelli S,Pinskerova V. "The shapes of the tibial and femoral articular surfaces in relation to tibiofemoral movement" .2002:11(3):pp172-186.
- [13] Moriguchi CS; Sato TO; Gil Coury HJC, "Ankle movements during normal gait evaluated by flexible electrogoniometer". *Rev. bras. fisioter*. vol.11 no.3 São Carlos May/June 2007.
- [14] Ramzi Sellaouti, Olivier Stasse, Shuuji Kajita, Kazuhito Yokoi and Abderrahmane Kheddar. "Faster and Smoother Walking of Humanoid HRP-2 with Passive Toe Joints" Post-doctoral Fellowship of Japan Society for Promotion of Science.
- [15] janne Heikkila and Olli Silve'n "A four step camera calibration procedure with implicit image correction". 13th International conference of pattern recognition. Vienna,Austria.
- [16] Kadaba MP, et al. *J. Orthop. Res* 7, 383-392, 1990

- [17] Jean Yves Bouguet, *The complete camera calibration Tool Box for Matlab* .computer vision research group Dep.of Electrical Engineering,California Institute of Technology.
- [18] National Institute of rehabilitation for disabled persons “National Institutes of Health Publication No. 93-3 509, March 1999.”
- [19] Kane TR, Levinson DA. *Dynamics: theory and application*. McGraw-Hill; New York, NY: 1985.
- [20] Arnold AS, Salinas S, Asakawa DJ, Delp SL. Accuracy of muscle moment arms estimated from MRI-based musculoskeletal models of the lower extremity. *Comput Aided Surg*. 2000; 5:108–119.
- [21] Maridy M. Troy, Hollye K. Moss , *Influence of step distance on knee flexion angle and cost during step exercise* ,Clinical Kinesiology: Journal of the American Kinesiotherapy Association , Fall, 2007.
- [22] Hoose, R. A. A., R. A. Lehnhard, T. Dewhurst, P. Pratt, and K. V. Hoose. Kinematic analysis of knee flexion during aerobic bench stepping and its clinical relevance. *Clin Kinesiol* 53:52-57, 1998.
- [23] Goodfellow, J., Hungerford, D. S. and C. Woods. Patello-femoral joint mechanics and pathology: 2. Chondromalacia Patellae. *J Bone Joint Surg* 58-B, 291-299, 1976.
- [24] Saunders JB, Inman VT, Eberhart HD. The major determinants in normal and pathological gait. *J Bone Joint Surg Am* 1953; 35-A:543-58.
- [25] Moseley L, Smith R, Hunt A, Gant R. Three-dimensional kinematics of the rearfoot during the stance phase of walking in normal young adult males. *Clinical Bio mechanics*. 1996;11:39-45.
- [26] Isacson J, Gransberg L, Knutsson E. Three-dimensional electrogoniometric gait recording. *Journal of Biomechanics*. 1986;19:627-35.

