

REFERENCES

- [1] H. N. Chanakya, H. K. Khuntia, N. Mukherjee, R. Aniruddha, J. R. Mudakavi, and P. Thimmaraju, “The physicochemical characteristics and anaerobic degradability of desiccated coconut industry waste water,” *Environmental Monitoring and Assessment*, vol. 187, no. 12, Nov. 2015, doi: <https://doi.org/10.1007/s10661-015-4991-7>.
- [2] B. Samarasiri, P. A. D. Mihiranga, and P. Rathnasiri, “Effect of lipid inhibition in anaerobic wastewater treatment: A case study using desiccated coconut wastewater,” doi: 10.13140/RG.2.2.36760.80646.
- [3] S. Kumar, G. Senanayake, C. Visvanathan, and B. Basu, “Desiccated coconut industry of Sri Lanka: opportunities for energy efficiency and environmental protection,” *Energy Conversion and Management*, vol. 44, no. 13, pp. 2205–2215, Aug. 2003, doi: [https://doi.org/10.1016/s0196-8904\(02\)00235-2](https://doi.org/10.1016/s0196-8904(02)00235-2).
- [4] B. K. T. Samarasiri, “Investigation on effect of lipid inhibition in anaerobic digestion of wastewater generated in desiccated coconut processing plants,” MSc theses, Department of Chemical and Process Engineering, University of Moratuwa, Sri Lanka, 2018.
- [5] H. V. A. Nandana and D. R. I. S. Werellagama, “Wastewater treatment in desiccated coconut industry,” *Proceedings of International Forestry and Environment Symposium*, vol. 0, no. 0, Aug. 2013, doi: <https://doi.org/10.31357/fesympo.v0i0.1613>.
- [6] H. N. Chanakya, H. K. Khuntia, N. Mukherjee, R. Aniruddha, J. R. Mudakavi, and P. Thimmaraju, “The physicochemical characteristics and anaerobic degradability of desiccated coconut industry waste water,” *Environmental Monitoring and Assessment*, vol. 187, no. 12, Nov. 2015, doi: <https://doi.org/10.1007/s10661-015-4991-7>.
- [7] D. G. Cirne, X. Paloumet, L. Björnsson, M. M. Alves, and B. Mattiasson, “Anaerobic digestion of lipid-rich waste - Effects of lipid concentration,” *Renewable Energy*, vol. 32, no. 6, pp. 965–975, May 2007, doi: <https://doi.org/10.1016/j.renene.2006.04.003>.
- [8] D. J. Batstone et al., “The IWA Anaerobic Digestion Model No 1 (ADM1),” 2002.
- [9] R. Sandriaty, C. Priadi, S. Kurnianingsih, and A. Abdillah, “Potential of biogas production from anaerobic co-digestion of fat, oil and grease waste and food waste,” *E3S Web of Conferences*, vol. 67, p. 02047, 2018, doi: <https://doi.org/10.1051/e3sconf/20186702047>.
- [10] N. Rasit, A. Idris, R. Harun, and W. A. Wan Ab Karim Ghani, “Effects of lipid inhibition on biogas production of anaerobic digestion from oily effluents and sludges: An overview,” *Renewable and Sustainable Energy Reviews*, vol. 45, pp. 351–358, May 2015, doi: <https://doi.org/10.1016/j.rser.2015.01.066>.

- [11] K. B. Chipasa and K. Mędrzycka, "Behavior of lipids in biological wastewater treatment processes," *Journal of Industrial Microbiology & Biotechnology*, vol. 33, no. 8, pp. 635–645, Feb. 2006, doi: <https://doi.org/10.1007/s10295-006-0099-y>.
- [12] K. Hanaki, T. Matsuo, and M. Nagase, "Mechanism of inhibition caused by long-chain fatty acids in anaerobic digestion process," *Biotechnology and Bioengineering*, vol. 23, no. 7, pp. 1591–1610, Jul. 1981, doi: <https://doi.org/10.1002/bit.260230717>.
- [13] K. T. Dasa, S. Y. Westman, R. Millati, M. N. Cahyanto, M. J. Taherzadeh, and C. Niklasson, "Inhibitory Effect of Long-Chain Fatty Acids on Biogas Production and the Protective Effect of Membrane Bioreactor," *BioMed Research International*, vol. 2016, p. 7263974, 2016, doi: <https://doi.org/10.1155/2016/7263974>.
- [14] A. Alhraishawi and S. Aslan, "Effect of Lipid Content on Anaerobic Digestion Process and Microbial Community: Review Study," *European Scientific Journal ESJ*, vol. 8, Aug. 2022, doi: <https://doi.org/10.19044/esipreprint.8.2022.p197>.
- [15] K. Dissanayake and P. G. Rathnasiri, "Effect of Lipid Inhibition in Batch Anaerobic Digesters Treating Desiccated Coconut Wastewater," *Annual sess. Of IESL*, pp. 467-475, 2019.
- [16] J. Ma et al., "Mechanism, kinetics and microbiology of inhibition caused by long-chain fatty acids in anaerobic digestion of algal biomass," *Biotechnology for Biofuels*, vol. 8, no. 1, Sep. 2015, doi: <https://doi.org/10.1186/s13068-015-0322-z>.
- [17] Ž. Zonta, M. M. Alves, X. Flotats, and J. Palatsi, "Modelling inhibitory effects of long chain fatty acids in the anaerobic digestion process," *Water Research*, vol. 47, no. 3, pp. 1369–1380, Mar. 2013, doi: <https://doi.org/10.1016/j.watres.2012.12.007>.
- [18] I. Angelidaki, L. Ellegaard, and B. K. Ahring, "A comprehensive model of anaerobic bioconversion of complex substrates to biogas," *Biotechnology and Bioengineering*, vol. 63, no. 3, pp. 363–372, May 1999, doi: [https://doi.org/10.1002/\(sici\)1097-0290\(19990505\)63:3%3C363::aid-bit13%3E3.0.co;2-z](https://doi.org/10.1002/(sici)1097-0290(19990505)63:3%3C363::aid-bit13%3E3.0.co;2-z).
- [19] E. Salminen, J. Rintala, L. Ya. Lokshina, and V. A. Vavilin, "Anaerobic batch degradation of solid poultry slaughterhouse waste," *Water Science and Technology*, vol. 41, no. 3, pp. 33–41, Feb. 2000, doi: <https://doi.org/10.2166/wst.2000.0053>.
- [20] L. Y. Lokshina, V. A. Vavilin, E. Salminen, and J. Rintala, "Modeling of Anaerobic Degradation of Solid Slaughterhouse Waste: Inhibition Effects of Long-Chain Fatty Acids or Ammonia," *Applied biochemistry and biotechnology*, vol. 109, no. 1–3, pp. 15–32, Jan. 2003, doi: <https://doi.org/10.1385/abab:109:1-3:15>.

- [21] H. N. Gavala, I. V. Skiadas, B. K. Ahring, and G. Lyberatos, "Thermophilic anaerobic fermentation of olive pulp for hydrogen and methane production: modelling of the anaerobic digestion process," *Water Science and Technology*, vol. 53, no. 8, pp. 271–279, Apr. 2006, doi: <https://doi.org/10.2166/wst.2006.258>.
- [22] J. Palatsi et al., "Long-chain fatty acids inhibition and adaptation process in anaerobic thermophilic digestion: Batch tests, microbial community structure and mathematical modelling," *Bioresource Technology*, vol. 101, no. 7, pp. 2243–2251, Apr. 2010, doi: <https://doi.org/10.1016/j.biortech.2009.11.069>.
- [23] A.B.G. Valladão, A. G. Torres, D. M. G. Freire, and M. C. Cammarota, "Profiles of fatty acids and triacylglycerols and their influence on the anaerobic biodegradability of effluents from poultry slaughterhouse," *Bioresource Technology*, vol. 102, no. 14, pp. 7043–7050, Apr. 2011, doi: <https://doi.org/10.1016/j.biortech.2011.04.037>.
- [24] B. K. McCabe, I. Hamawand, and C. Baillie, "Investigating wastewater modelling as a tool to predict anaerobic decomposition and biogas yield of abattoir effluent," *Journal of Environmental Chemical Engineering*, vol. 1, no. 4, pp. 1375–1379, Dec. 2013, doi: <https://doi.org/10.1016/j.jece.2013.07.015>.
- [25] R. Petruy and G. Lettinga, "Digestion of a milk-fat emulsion," *Bioresource Technology*, vol. 61, no. 2, pp. 141–149, Aug. 1997, doi: [https://doi.org/10.1016/s0960-8524\(97\)00042-4](https://doi.org/10.1016/s0960-8524(97)00042-4).