

References

1. Agustín Martínez, G., Bertin, L., Scoma, A., Rebecchi, S., Braunegg, G., & Fava, F. (2015). Production of polyhydroxyalkanoates from dephenolised and fermented olive mill wastewaters by employing a pure culture of *Cupriavidus necator*. *Biochemical Engineering Journal*, *97*, 92–100. <https://doi.org/10.1016/j.bej.2015.02.015>
2. Ajila, C. M., & Prasada Rao, U. J. S. (2013). Mango peel dietary fibre: Composition and associated bound phenolics. *Journal of Functional Foods*, *5*(1), 444–450. <https://doi.org/10.1016/j.jff.2012.11.017>
3. Amini, M., Yousefi-Massumabad, H., Younesi, H., Abyar, H., & Bahramifar, N. (2020). Production of the polyhydroxyalkanoate biopolymer by *Cupriavidus necator* using beer brewery wastewater containing maltose as a primary carbon source. *Journal of Environmental Chemical Engineering*, *8*(1). <https://doi.org/10.1016/j.jece.2019.103588>
4. Ammar, E. M., El-Sheshtawy, H. S., El-Shatoury, E. H., & Amer, S. K. (2021). Green synthesis of polyhydroxyalkanoate polymer by *Bacillus iocasae*. *Polymer International*, *70*(10), 1478–1485. <https://doi.org/10.1002/pi.6219>
5. Anjali, M., Sukumar, C., Kanakalakshmi, A., & Shanthi, K. (2014). Enhancement of growth and production of polyhydroxyalkanoates by *Bacillus subtilis* from agro-industrial waste as carbon substrates. *Composite Interfaces*, *21*(2), 111–119. <https://doi.org/10.1080/15685543.2013.834200>
6. Balakrishna Pillai, A., Jaya Kumar, A., Thulasi, K., & Kumarapillai, H. (2017). Evaluation of short-chain-length polyhydroxyalkanoate accumulation in *Bacillus aryabhatai*. *Brazilian Journal of Microbiology*, *48*(3), 451–460. <https://doi.org/10.1016/j.bjm.2017.01.005>
7. Bartolomé, A. P., Rupprez, P., & Fhster, C. (1995). Pineapple fruit: morphological characteristics, chemical composition and sensory analysis of Red Spanish and Smooth Cayenne cultivars. In *Food Chemistry* (Vol. 53).
8. Benesova, P., Kucera, D., Marova, I., & Obruca, S. (2017). Chicken feather hydrolysate as an inexpensive complex nitrogen source for PHA production by *Cupriavidus necator* on waste frying oils. *Letters in Applied Microbiology*, *65*(2), 182–188. <https://doi.org/10.1111/lam.12762>

9. Brojanigo, S., Parro, E., Cazzorla, T., Favaro, L., Basaglia, M., & Casella, S. (2020). Conversion of starchy waste streams into polyhydroxyalkanoates using *Cupriavidus necator* DSM 545. *Polymers*, *12*(7). <https://doi.org/10.3390/polym12071496>
10. Bugnicourt, E., Cinelli, P., Lazzeri, A., & Alvarez, V. (2014). Polyhydroxyalkanoate (PHA): Review of synthesis, characteristics, processing and potential applications in packaging. *Express Polymer Letters*, *8*(11), 791–808. <https://doi.org/10.3144/expresspolymlett.2014.82>
11. Cavaliere, C., Montone, C. M., Capriotti, A. L., la Barbera, G., Piovesana, S., Rotatori, M., Valentino, F., & Laganà, A. (2018). Extraction of polycyclic aromatic hydrocarbons from polyhydroxyalkanoates before gas chromatography/mass spectrometry analysis. *Talanta*, *188*, 671–675. <https://doi.org/10.1016/j.talanta.2018.06.038>
12. Dalton, B., Bhagabati, P., de Micco, J., Padamati, R. B., & O'Connor, K. (2022). A Review on Biological Synthesis of the Biodegradable Polymers Polyhydroxyalkanoates and the Development of Multiple Applications. In *Catalysts* (Vol. 12, Issue 3). MDPI. <https://doi.org/10.3390/catal12030319>
13. de Silva, L., Doremus, J., & Taylor, R. L. C. (2021). *The Plastic Economy A review of the positive and negative impacts of plastic and its alternatives*. <https://ssrn.com/abstract=3859528>
14. de Smet, M. J., Eggink, G., Witholt, B., Kingma, J., & Wynberg, H. (1983). Characterization of Intracellular Inclusions Formed by *Pseudomonas oleovorans* During Growth on Octane. In *JOURNAL OF BACTERIOLOGY* (Vol. 154, Issue 2). <https://journals.asm.org/journal/jb>
15. de Souza Reis, G. A., Michels, M. H. A., Fajardo, G. L., Lamot, I., & de Best, J. H. (2020). Optimization of green extraction and purification of PHA produced by mixed microbial cultures from sludge. *Water (Switzerland)*, *12*(4). <https://doi.org/10.3390/W12041185>
16. Din, M. I., Ghaffar, T., Najeeb, J., Hussain, Z., Khalid, R., & Zahid, H. (2020). Potential perspectives of biodegradable plastics for food packaging application-review of properties and recent developments. In *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment* (Vol. 37, Issue 4, pp. 665–680). Taylor and Francis Ltd. <https://doi.org/10.1080/19440049.2020.1718219>

17. dos Santos, A. J., Oliveira Dalla Valentina, L. V., Hidalgo Schulz, A. A., & Tomaz Duarte, M. A. (2017). From Obtaining to Degradation of PHB:Material Properties. Part I. *Ingeniería y Ciencia*, 13(26), 269–298. <https://doi.org/10.17230/ingciencia.13.26.10>
18. Dunuweera, A. N., Nikagolla, D. N., & Ranganathan, K. (2021). Fruit Waste Substrates to Produce Single-Cell Proteins as Alternative Human Food Supplements and Animal Feeds Using Baker's Yeast (*Saccharomyces cerevisiae*). *Journal of Food Quality*, 2021. <https://doi.org/10.1155/2021/9932762>
19. Duque, A. F., Oliveira, C. S. S., Carmo, I. T. D., Gouveia, A. R., Pardelha, F., Ramos, A. M., & Reis, M. A. M. (2014). Response of a three-stage process for PHA production by mixed microbial cultures to feedstock shift: Impact on polymer composition. *New Biotechnology*, 31(4), 276–288. <https://doi.org/10.1016/j.nbt.2013.10.010>
20. Ebu, S. M., & Ray, L. (2021). *Polyhydroxyalkanoate Recovery From Newly Screened Bacillus Sp. LPPI-81 Using Various Methods of Extraction From Loktak Lake Sediment Sample*. <https://doi.org/10.21203/rs.3.rs-1060626/v1>
21. (EDB). (2019). *INDUSTRY CAPABILITY REPORT FRESH FRUIT & VEGETABLE*,.
22. Elbanna, K., Lütke-Eversloh, T., Jendrossek, D., Luftmann, H., & Steinbüchel, A. (2004). Studies on the biodegradability of polythioester copolymers and homopolymers by polyhydroxyalkanoate (PHA)-degrading bacteria and PHA depolymerases. *Archives of Microbiology*, 182(2–3), 212–225. <https://doi.org/10.1007/s00203-004-0715-z>
23. Fang, F., Xu, R. Z., Huang, Y. Q., Wang, S. N., Zhang, L. L., Dong, J. Y., Xie, W. M., Chen, X., & Cao, J. S. (2019). Production of polyhydroxyalkanoates and enrichment of associated microbes in bioreactors fed with rice winery wastewater at various organic loading rates. *Bioresource Technology*, 292. <https://doi.org/10.1016/j.biortech.2019.121978>
24. Flores-Sánchez, A., López-Cuellar, M. D. R., Pérez-Guevara, F., Figueroa López, U., Martín-Bufájer, J. M., & Vergara-Porras, B. (2017). Synthesis of Poly-(R-hydroxyalkanoates) by *Cupriavidus necator* ATCC 17699 Using Mexican Avocado (*Persea americana*) Oil as a Carbon Source. *International Journal of Polymer Science*, 2017. <https://doi.org/10.1155/2017/6942950>
25. Gobi, K., & Vadivelu, V. M. (2015). Polyhydroxyalkanoate recovery and effect of in situ extracellular polymeric substances removal from aerobic granules. *Bioresource Technology*, 189, 169–176. <https://doi.org/10.1016/j.biortech.2015.04.023>

26. Gomaa, E. Z. (2014). BRAZILIAN ARCHIVES OF BIOLOGY AND TECHNOLOGY Production of Polyhydroxyalkanoates (PHAs) By *Bacillus subtilis* and *Escherichia coli* Grown on Cane Molasses Fortified with Ethanol. *Arch. Biol. Technol.* v, 57(1), 145–154.
27. Gusakov, A. v., Kondratyeva, E. G., & Sinitsyn, A. P. (2011). Comparison of Two Methods for Assaying Reducing Sugars in the Determination of Carbohydrase Activities. *International Journal of Analytical Chemistry*, 2011, 1–4. <https://doi.org/10.1155/2011/283658>
28. Hairudin, N. H. B. M., Ganesan, S., & Sudesh, K. (2021). Revalorization of adsorbed residual oil in spent bleaching clay as a sole carbon source for polyhydroxyalkanoate (PHA) accumulation in *Cupriavidus necator* Re2058/pCB113. *Polymer Journal*, 53(1), 169–178. <https://doi.org/10.1038/s41428-020-00418-2>
29. Hazer, B., & Steinbüchel, A. (2007). Increased diversification of polyhydroxyalkanoates by modification reactions for industrial and medical applications. In *Applied Microbiology and Biotechnology* (Vol. 74, Issue 1, pp. 1–12). <https://doi.org/10.1007/s00253-006-0732-8>
30. Hoben, H. J., & Somasegaran, P. (1982). Comparison of the Pour, Spread, and Drop Plate Methods for Enumeration of *Rhizobium* spp. in Inoculants Made from Presterilized Peatt. In *APPLIED AND ENVIRONMENTAL MICROBIOLOGY* (Vol. 44, Issue 5). <https://journals.asm.org/journal/aem>
31. Huang, J.-C., Shew, A. S., & Wang, M.-S. (1990). *Biodegradable Plastics: A Review*.
32. Hu, R., Lin, L., Liu, T., Ouyang, P., He, B., & Liu, S. (2008). Reducing sugar content in hemicellulose hydrolysate by DNS method: A revisit. *Journal of Biobased Materials and Bioenergy*, 2(2), 156–161. <https://doi.org/10.1166/jbmb.2008.306>
33. Kaniuk, Ł., & Stachewicz, U. (2021). Development and Advantages of Biodegradable PHA Polymers Based on Electrospun PHBV Fibers for Tissue Engineering and Other Biomedical Applications. In *ACS Biomaterials Science and Engineering* (Vol. 7, Issue 12, pp. 5339–5362). American Chemical Society. <https://doi.org/10.1021/acsbiomaterials.1c00757>
34. Kemavongse, K., Prasertsan, P., Upaichit, A., & Methacanon, P. (2008). Poly- β -hydroxyalkanoate production by halotolerant *Rhodobacter sphaeroides* U7. *World Journal*

of *Microbiology and Biotechnology*, 24(10), 2073–2085. <https://doi.org/10.1007/s11274-008-9712-8>

35. Kim, S. W., Baek, Y. W., & An, Y. J. (2011). Assay-dependent effect of silver nanoparticles to *Escherichia coli* and *Bacillus subtilis*. *Applied Microbiology and Biotechnology*, 92(5), 1045–1052. <https://doi.org/10.1007/s00253-011-3611-x>

36. Koller, M. (2018a). Biodegradable and biocompatible polyhydroxy-alkanoates (PHA): Auspicious microbial macromolecules for pharmaceutical and therapeutic applications. In *Molecules* (Vol. 23, Issue 2). MDPI AG. <https://doi.org/10.3390/molecules23020362>

37. Koller, M. (2018b). A review on established and emerging fermentation schemes for microbial production of polyhydroxyalkanoate (PHA) biopolyesters. In *Fermentation* (Vol. 4, Issue 2). MDPI AG. <https://doi.org/10.3390/fermentation4020030>

38. Kovalcik, A., Pernicova, I., Obruca, S., Szotkowski, M., Enev, V., Kalina, M., & Marova, I. (2020a). Grape winery waste as a promising feedstock for the production of polyhydroxyalkanoates and other value-added products. *Food and Bioproducts Processing*, 124, 1–10. <https://doi.org/10.1016/j.fbp.2020.08.003>

39. Kovalcik, A., Pernicova, I., Obruca, S., Szotkowski, M., Enev, V., Kalina, M., & Marova, I. (2020b). Grape winery waste as a promising feedstock for the production of polyhydroxyalkanoates and other value-added products. *Food and Bioproducts Processing*, 124, 1–10. <https://doi.org/10.1016/j.fbp.2020.08.003>

40. Kumari, B. (2014). Studies on pectinase production by *Bacillus subtilis* using agro-industrial wastes Pharmacology View project Analytical method development View project. In *Article in Research Journal of Pharmaceutical, Biological and Chemical Sciences*. <https://www.researchgate.net/publication/286658976>

41. Lebaka, V. R., Wee, Y. J., Ye, W., & Korivi, M. (2021). Nutritional composition and bioactive compounds in three different parts of mango fruit. In *International Journal of Environmental Research and Public Health* (Vol. 18, Issue 2, pp. 1–20). MDPI AG. <https://doi.org/10.3390/ijerph18020741>

42. Levett, I., Birkett, G., Davies, N., Bell, A., Langford, A., Laycock, B., Lant, P., & Pratt, S. (2016). Techno-economic assessment of poly-3-hydroxybutyrate (PHB) production from methane - The case for thermophilic bioprocessing. *Journal of Environmental Chemical Engineering*, 4(4), 3724–3733. <https://doi.org/10.1016/j.jece.2016.07.033>

43. Lyla, A. (2020). Theoretical study on bio-based material. *Malaya Journal of Matematik*, *S(2)*. <https://doi.org/10.26637/MJM0S20/0324>
44. Mahato, R. P., Kumar, S., & Singh, P. (2021a). Optimization of Growth Conditions to Produce Sustainable Polyhydroxyalkanoate Bioplastic by *Pseudomonas aeruginosa* EO1. *Frontiers in Microbiology*, *12*. <https://doi.org/10.3389/fmicb.2021.711588>
45. Mahato, R. P., Kumar, S., & Singh, P. (2021b). Optimization of Growth Conditions to Produce Sustainable Polyhydroxyalkanoate Bioplastic by *Pseudomonas aeruginosa* EO1. *Frontiers in Microbiology*, *12*. <https://doi.org/10.3389/fmicb.2021.711588>
46. Maier, R. M. (2000). *37 3.1 Growth in Pure Culture in a Flask 3.1.1 The Lag Phase 3.1.2 The Exponential Phase 3.1.3 The Stationary Phase 3.1.4 The Death Phase 3.1.5 Effect of Substrate Concentration on Growth Bacterial Growth*.
47. Maldonado-Celis, M. E., Yahia, E. M., Bedoya, R., Landázuri, P., Loango, N., Aguillón, J., Restrepo, B., & Guerrero Ospina, J. C. (2019). Chemical Composition of Mango (*Mangifera indica* L.) Fruit: Nutritional and Phytochemical Compounds. In *Frontiers in Plant Science* (Vol. 10). Frontiers Media S.A. <https://doi.org/10.3389/fpls.2019.01073>
48. Marfo, E. K., Oke, O. L., & Afolabi, O. A. (1986). Chemical Composition of Papaya (*Carica papaya*) Seeds. In *Food Chemistry* (Vol. 22).
49. Markl, E. (2018). PHB - Bio Based and Biodegradable Replacement for PP: A Review. *Novel Techniques in Nutrition & Food Science*, *2(5)*. <https://doi.org/10.31031/ntnf.2018.02.000546>
50. Marsden, W. L., Gray, P. P., Nippard, G. J., & Quinlan, M. R. (1982). Evaluation of the DNS Method for Analysing Lignocellulosic Hydrolysates. In *J. Chem. Tech. Biolechnol* (Vol. 32).
51. Mitra, R., Xu, T., Xiang, H., & Han, J. (2020). Current developments on polyhydroxyalkanoates synthesis by using halophiles as a promising cell factory. In *Microbial Cell Factories* (Vol. 19, Issue 1). BioMed Central Ltd. <https://doi.org/10.1186/s12934-020-01342-z>
52. Mohapatra, S., Maity, S., Dash, H. R., Das, S., Pattnaik, S., Rath, C. C., & Samantaray, D. (2017). Bacillus and biopolymer: Prospects and challenges. In *Biochemistry and Biophysics Reports* (Vol. 12, pp. 206–213). Elsevier B.V. <https://doi.org/10.1016/j.bbrep.2017.10.001>

53. Mohapatra, S., Mohanta, P. R., Sarkar, B., Daware, A., Kumar, C., & Samantaray, D. P. (2017). Production of Polyhydroxyalkanoates (PHAs) by Bacillus Strain Isolated from Waste Water and Its Biochemical Characterization. *Proceedings of the National Academy of Sciences India Section B - Biological Sciences*, 87(2), 459–466. <https://doi.org/10.1007/s40011-015-0626-6>
54. Morais, D. R., Rotta, E. M., Sargi, S. C., Bonafe, E. G., Suzuki, R. M., Souza, N. E., Matsushita, M., & Visentainer, J. v. (2017). Proximate composition, mineral contents and fatty acid composition of the different parts and dried peels of tropical fruits cultivated in Brazil. *Journal of the Brazilian Chemical Society*, 28(2), 308–318. <https://doi.org/10.5935/0103-5053.20160178>
55. Mozumder, M. S. I., de Wever, H., Volcke, E. I. P., & Garcia-Gonzalez, L. (2014). A robust fed-batch feeding strategy independent of the carbon source for optimal polyhydroxybutyrate production. *Process Biochemistry*, 49(3), 365–373. <https://doi.org/10.1016/j.procbio.2013.12.004>
56. Muhammadi, Shabina, Afzal, M., & Hameed, S. (2015). Bacterial polyhydroxyalkanoates-eco-friendly next generation plastic: Production, biocompatibility, biodegradation, physical properties and applications. *Green Chemistry Letters and Reviews*, 8(3–4), 56–77. <https://doi.org/10.1080/17518253.2015.1109715>
57. Nambu, Y., Ishii-Hyakutake, M., Harada, K., Mizuno, S., & Tsuge, T. (2020). Expanded amino acid sequence of the PhaC box in the active center of polyhydroxyalkanoate synthases. *FEBS Letters*, 594(4), 710–716. <https://doi.org/10.1002/1873-3468.13651>
58. Naser, A. Z., Deiab, I., & Darras, B. M. (2021). Poly(lactic acid) (PLA) and polyhydroxyalkanoates (PHAs), green alternatives to petroleum-based plastics: a review. In *RSC Advances* (Vol. 11, Issue 28, pp. 17151–17196). Royal Society of Chemistry. <https://doi.org/10.1039/d1ra02390j>
59. Nyssa Nair, Steven Pratt, Paul Lant, & Bronwyn Laycock. (2022). Bioplastics Promise. *The Chemical Engineer*.
60. Obruca, S., Marova, I., Snajdar, O., Mravcova, L., & Svoboda, Z. (2010). Production of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) by *Cupriavidus necator* from waste rapeseed oil using propanol as a precursor of 3-hydroxyvalerate. *Biotechnology Letters*, 32(12), 1925–1932. <https://doi.org/10.1007/s10529-010-0376-8>

61. Ojimelukwe, P., Nwofia, E., Ojimelukwe, P., & Eji, C. (2012). Chemical composition of leaves, fruit pulp and seeds in some *Carica papaya* (L) morphotypes Production and evaluation of teas from plant products, for the management of type 2 diabetes View project Isolation, modification and evaluation of icacina trichantha tuber View project Chemical composition of leaves, fruit pulp and seeds in some *Carica papaya* (L) morphotypes. In *Int. J. Med. Arom. Plants* (Vol. 2, Issue 1). <http://www.openaccessscience.com>
62. Olayinka, B. U. (2018). *Proximate and Chemical Compositions of Watermelon (Citrullus lanatus (Thunb.) Matsum and Nakai cv Red and Cucumber (Cucumis sativus L. cv Pipino)*.
63. Park, D. H., & Kim, B. S. (2011). Production of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-4-hydroxybutyrate) by *Ralstonia eutropha* from soybean oil. *New Biotechnology*, 28(6), 719–724. <https://doi.org/10.1016/j.nbt.2011.01.007>
64. Petrovova, M., Tkadlec, J., Dvoracek, L., Streitova, E., & Licha, I. (2014). NAD(P)H-hydrate dehydratase- A metabolic repair enzyme and its role in *Bacillus subtilis* stress adaptation. *PLoS ONE*, 9(11). <https://doi.org/10.1371/journal.pone.0112590>
65. *Plastic Waste Management - Sri Lanka*. (2021). <https://www.unep.org/interactive/beat-plastic-pollution/>
66. Prasertsung, I., Aroonraj, K., Kamwilaisak, K., Saito, N., & Damrongsakkul, S. (2019). Production of reducing sugar from cassava starch waste (CSW) using solution plasma process (SPP). *Carbohydrate Polymers*, 205, 472–479. <https://doi.org/10.1016/j.carbpol.2018.10.090>
67. Purama, R. K., Al-Sabahi, J. N., & Sudesh, K. (2018). Evaluation of date seed oil and date molasses as novel carbon sources for the production of poly(3Hydroxybutyrate-co-3Hydroxyhexanoate) by *Cupriavidus necator* H16 Re 2058/pCB113. *Industrial Crops and Products*, 119, 83–92. <https://doi.org/10.1016/j.indcrop.2018.04.013>
68. Rayasam, V., Chavan, P., & Kumar, T. (2020). Polyhydroxyalkanoate synthesis by bacteria isolated from landfill and ETP with pomegranate peels as carbon source. *Archives of Microbiology*, 202(10), 2799–2808. <https://doi.org/10.1007/s00203-020-01995-9>
69. Reddy, C., Ghai, R., & Kalia, V. (2003). *Polyhydroxyalkanoates: an overview*.
70. Rico, X., Gullón, B., Alonso, J. L., & Yáñez, R. (2020). Recovery of high value-added compounds from pineapple, melon, watermelon and pumpkin processing by-products: An

- overview. In *Food Research International* (Vol. 132). Elsevier Ltd. <https://doi.org/10.1016/j.foodres.2020.109086>
71. Samarasinghe, K., Pawan Kumar, S., & Visvanathan, C. (2021). Evaluation of circular economy potential of plastic waste in Sri Lanka. *Environmental Quality Management*, 31(1), 99–107. <https://doi.org/10.1002/tqem.21732>
72. Satoh, H., Iwamoto, Y., Mino, T., & Matsuo, T. (1998). Activated sludge as a possible source of biodegradable plastic. *Water Science and Technology*, 38(2 pt 2), 103–109. [https://doi.org/10.1016/S0273-1223\(98\)00435-1](https://doi.org/10.1016/S0273-1223(98)00435-1)
73. Shah K R. (2011). *ftir-with-cover-page-v2*.
74. Shah, S., & Kumar, A. (2020). Polyhydroxyalkanoates: Advances in the synthesis of sustainable bio-plastics. *European Journal of Environmental Sciences*, 10(2), 76–88. <https://doi.org/10.14712/23361964.2020.9>
75. Stavroula, K., Simos, M., & Katherine-Joanne, H. (2020). Polyhydroxyalkanoates (PHAs) from Household Food Waste: Research Over the Last Decade. *International Journal of Biotechnology and Bioengineering*, 6(2), 26–36.
76. Stevenson, K., McVey, A. F., Clark, I. B. N., Swain, P. S., & Pilizota, T. (2016). General calibration of microbial growth in microplate readers. *Scientific Reports*, 6. <https://doi.org/10.1038/srep38828>
77. Sukruansuwan, V., & Napathorn, S. C. (2018). Use of agro-industrial residue from the canned pineapple industry for polyhydroxybutyrate production by *Cupriavidus necator* strain A-04. *Biotechnology for Biofuels*, 11(1). <https://doi.org/10.1186/s13068-018-1207-8>
78. Thiviya, P., Gamage, A., Kapilan, R., Merah, O., & Madhujith, T. (2022). Single Cell Protein Production Using Different Fruit Waste: A Review. *Separations*, 9(7), 178. <https://doi.org/10.3390/separations9070178>
79. Thomas, P., Sekhar, A. C., Upreti, R., Mujawar, M. M., & Pasha, S. S. (2015). Optimization of single plate-serial dilution spotting (SP-SDS) with sample anchoring as an assured method for bacterial and yeast cfu enumeration and single colony isolation from diverse samples. *Biotechnology Reports*, 8, 45–55. <https://doi.org/10.1016/j.btre.2015.08.003>

80. Thomson, N., Summers, D., & Sivaniah, E. (2010). Synthesis, properties and uses of bacterial storage lipid granules as naturally occurring nanoparticles. In *Soft Matter* (Vol. 6, Issue 17, pp. 4045–4057). <https://doi.org/10.1039/b927559b>
81. Umesh, M., Priyanka, K., Thazeem, B., & Preethi, K. (2018). Biogenic PHA nanoparticle synthesis and characterization from *Bacillus subtilis* NCDC0671 using orange peel medium. *International Journal of Polymeric Materials and Polymeric Biomaterials*, 67(17), 996–1004. <https://doi.org/10.1080/00914037.2017.1417284>
82. Verlinden, R. A. J., Hill, D. J., Kenward, M. A., Williams, C. D., & Radecka, I. (2007). Bacterial synthesis of biodegradable polyhydroxyalkanoates. In *Journal of Applied Microbiology* (Vol. 102, Issue 6, pp. 1437–1449). <https://doi.org/10.1111/j.1365-2672.2007.03335.x>
83. Wang, Q., Xia, Y., Chen, Q., & Qi, Q. (2012). Incremental truncation of PHA synthases results in altered product specificity. *Enzyme and Microbial Technology*, 50(6–7), 293–297. <https://doi.org/10.1016/j.enzmictec.2012.02.003>
84. Xiao, N., & Jiao, N. (2011). Formation of polyhydroxyalkanoate in aerobic anoxygenic phototrophic bacteria and its relationship to carbon source and light availability. *Applied and Environmental Microbiology*, 77(21), 7445–7450. <https://doi.org/10.1128/AEM.05955-11>
85. Yan, Z., Sousa-Gallagher, M. J., & Oliveira, F. A. R. (2008). Shrinkage and porosity of banana, pineapple and mango slices during air-drying. *Journal of Food Engineering*, 84(3), 430–440. <https://doi.org/10.1016/j.jfoodeng.2007.06.004>
86. Yu, C., Fan, L., Gao, J., Wang, M., Wu, Q., Tang, J., Li, Y., & Chen, J. (2015). The platelet-activating factor acetylhydrolase gene derived from *Trichoderma harzianum* induces maize resistance to *Curvularia lunata* through the jasmonic acid signaling pathway. *Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes*, 50(10), 708–717. <https://doi.org/10.1080/03601234.2015.1048104>
87. Zainab-L, I., Ng, W., & Sudesh, K. (2022). Potential of mealworms used in polyhydroxyalkanoate/bioplastic recovery as red hybrid tilapia (*Oreochromis* sp.) feed ingredient. *Scientific Reports*, 12(1). <https://doi.org/10.1038/s41598-022-13429-1>
88. Zaki, Dr. N. H. (2018). Biodegradable plastic production by *Bacillus* spp. isolated from agricultural wastes and genetic analysis of PHA synthesis. *Al-Mustansiriyah Journal of Science*, 29(1), 67–74. <https://doi.org/10.23851/mjs.v29i1.115>